

AD-A090 797

STANFORD UNIV CA SYSTEMS OPTIMIZATION LAB

F/G 12/1

A CONSTRUCTIVE PROOF OF TUCKER'S COMBINATORIAL LEMMA.(U)

JUN 80 R M FREUND, M J TODD

DAA629-78-6-0026

UNCLASSIFIED

SOL-80-12

ARO-15254.10-M

NL

1 OF 1  
AD A  
7990797



END  
DATE  
FILMED  
11-80  
DTIC

ARO 15254.10-M



Systems  
Optimization  
Laboratory

LEVEL II

(11)

AD A090797

DDC FILE COPY.

DTIC  
ELECTE  
OCT 23 1980  
S D  
E

DISTRIBUTION STATEMENT A

Approved for public release  
Distribution Unlimited

Department of Operations Research  
Stanford University  
Stanford, CA 94305

80 10 20 019

**LEVEL II**

(11)

**SYSTEMS OPTIMIZATION LABORATORY  
DEPARTMENT OF OPERATIONS RESEARCH  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA 94305**

**A CONSTRUCTIVE PROOF OF TUCKER'S  
COMBINATORIAL LEMMA**

by

**Robert A. Freund\***  
and **Michael J. Todd\*\***

**TECHNICAL REPORT SOL 80-12  
June 1980**

\*Research and reproduction of this report were partially supported by the Department of Energy Contract DE-AC03-76-SF00326, PA No. DE-AT-03-76ER72018; the National Science Foundation Grants MCS79-03145 and SOC78-16811; U.S. Army Research Office contract DAAG-29-78-G-0026, at Stanford University.

\*\*Research of this report was partially supported by National Science Foundation Grant ENG76-08749 at Cornell University.

Reproduction in whole or in part is permitted for any purposes of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.

**DISTRIBUTION STATEMENT A**

Approved for public release;

## Abstract

Tucker's combinatorial lemma is concerned with certain labellings of the vertices of a triangulation of the  $n$ -ball. It can be used as a basis for the proof of antipodal-point theorems in the same way that Sperner's lemma yields Brouwer's theorem. Here we give a constructive proof, which thereby yields algorithms for antipodal-point problems. Our method is based on an algorithm of Reiser.

[illegible]

## Introduction

Let  $B^n$  denote the  $n$ -ball  $\{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$  where  $\|x\|$  is the  $\ell_1$ -norm  $\sum_i |x_i|$ , and let  $S^{n-1}$  denote its boundary  $\{x \in \mathbb{R}^n \mid \|x\| = 1\}$ . We will call special a centrally symmetric triangulation of  $B^n$  that refines the octahedral subdivision. The following result was proved for  $n = 2$  in [6]; for the general case, see [2, pages 134-141].

Tucker's Combinatorial Lemma. Let the vertices of a special triangulation  $T$  of  $B^n$  be assigned labels from  $\{\pm 1, \dots, \pm n\}$ . If antipodal vertices of  $T$  on  $S^{n-1}$  receive complementary labels (labels that sum to zero), then  $T$  contains a complementary 1-simplex (a 1-simplex whose vertices have complementary labels).

Tucker stated his lemma in a different form; the nonexistence of such a labelling with no complementary 1-simplex was asserted. (In [2] and an earlier abstract [7] a related positive assertion is given; however we have not been able to find a constructive proof of this lemma.) We will prove below that a complementary 1-simplex exists by devising an algorithm that will find one. The algorithm is based on a method of Reiser for the nonlinear complementarity problem [4].

Not only is Tucker's lemma stated in [6] in terms of nonexistence; his derivation of antipodal-point theorems from it was by contradiction. We briefly indicate below how constructive proofs of two such theorems follow from our algorithm.

The Borsuk-Ulam Theorem. If a continuous function maps  $S^n$  into  $\mathbb{R}^n$ , at least one pair of antipodal points is mapped into a single point.

Proof. Let the function be  $f: S^n \rightarrow \mathbb{R}^n$  and define  $g: B^n \rightarrow \mathbb{R}^n$  by  $g(x) = f(x, 1 - \|x\|) - f(-x, \|x\| - 1)$ ; note that  $g(-x) = -g(x)$  for  $x \in S^{n-1}$ .

Now for any special triangulation  $T$  of  $B^n$ , label vertex  $v + i$  ( $-i$ ) if  $|g_i(v)| = \max_j |g_j(v)|$  and  $g_i(v)$  is positive (negative) (if  $g(v) = 0$ , we are done). In case of ties, the least such index is chosen. This labelling satisfies the requirements of the lemma and hence  $T$  contains a complementary 1-simplex. Let  $x^*$  be any limit point of such complementary 1-simplices for a sequence of special triangulations whose meshes approach zero. A continuity argument implies that  $g(x^*) = 0$ , and hence  $(x^*, 1 - \|x^*\|)$  and its antipode are mapped by  $f$  into the same point.

The Lusternik-Schnirelmann Theorem. If  $S^n$  is covered by  $n+1$  closed sets, at least one of them contains a pair of antipodal points.

Proof. Let the sets be  $C_1, C_2, \dots, C_{n+1}$ . Label the vertices of any special triangulation  $T$  of  $B^n$  as follows. If  $v$  is such a vertex, let  $y = (v, 1 - \|v\|)$ . Determine the least  $i$  such that  $C_i$  contains  $y$  or  $-y$ . If  $C_i$  contains both  $y$  and  $-y$ , the theorem is proved. Also, if  $i = n+1$  then since neither  $y$  nor  $-y$  lie in  $C_j$ ,  $1 \leq j \leq n$ , both must lie in  $C_{n+1}$  and we are done. Hence assume that for each vertex  $v$ ,  $i$  is between 1 and  $n$  and  $C_i$  contains one of  $y, -y$ . Label  $v + i$  or  $-i$  accordingly. Again, the labelling satisfies the requirements of the lemma and a complementary 1-simplex exists. Similarly, a limiting argument (some complementary pair of labels occurs infinitely often; take any limit point of the corresponding complementary 1-simplices) proves the claim.

Meyerson and Wright [3] and Barany [1] have also given algorithms for the Borsuk-Ulam theorem. Both use vector-labelling, and their algorithms should be more efficient for practical problems; however no computationally tractable way is yet known to implement their techniques for dealing with the theoretical possibility of degeneracy. We note that in an algorithm it would be preferable

to use the  $n$ -ball induced by the  $\ell_\infty$ -norm,  $B_\infty^n = \{x \in \mathbb{R}^n \mid |x_i| \leq 1 \text{ all } i\}$ ; several efficient special triangulations of  $B_\infty^n$  exist, for example,  $K_1$  and  $J_1$  [5, Chapter III]. Figure 1 below shows  $B_\infty^2$  rather than  $B^2$  for this reason.

### The Algorithm

First we need some notation. We note by  $\text{sgn}(\lambda)$  the sign (0, +1 or -1) of any real number  $\lambda$ ; similarly, for a vector  $x = (x_i) \in \mathbb{R}^n$ ,  $\text{sgn}(x)$  is the vector  $(\text{sgn}(x_i))$ . For any sign vector  $s \in \mathbb{R}^n$  (i.e., each component  $s_i$  is 0, +1 or -1),  $C(s)$  denotes the closure of  $\{x \in \mathbb{R}^n \mid \text{sgn}(x) = s\}$ , i.e. the set of those  $x$  for which  $x_i$  is nonnegative, zero or nonpositive according as  $s_i$  is 1, 0, or -1, for each  $i$ . We call  $C(s)$  an orthant - actually it is an orthant of a coordinate subspace. Any special triangulation  $T$  induces triangulations of  $C(s) \cap B^n$  for each  $s$ . Let  $\sigma$  be a simplex of  $T$ ; then  $\text{sgn}(x)$  is the same for each  $x$  in the relative interior of  $\sigma$  -- we let  $\text{sgn}(\sigma)$  be this sign vector. Clearly,  $C(\text{sgn}(\sigma))$  is the smallest orthant containing  $\sigma$ . Since  $T$  is centrally symmetric, every simplex  $\sigma$  lying in  $S^{n-1}$  has an antipodal simplex  $-\sigma = \{-x \mid x \in \sigma\}$ .

Definition. For any sign vector  $s$ , a simplex  $\sigma \in T$  is  $s$ -labelled if, whenever  $s_i$  is nonzero,  $s_i i$  is a label of some vertex of  $\sigma$ . If  $\sigma$  is  $\text{sgn}(\sigma)$ -labelled, we say  $\sigma$  is completely labelled.

Note that the 0-simplex  $\{0\}$  is always completely labelled by default since its sign vector is zero. Also, if  $\sigma \subseteq S^{n-1}$  is completely labelled, so is its antipodal simplex.

The algorithm proceeds by tracing a path in a graph  $G$  whose nodes are completely labelled simplices until it finds a complementary 1-simplex. The graph is given by the following:

Definition. Two completely labelled simplices  $\sigma$  and  $\tau$  are adjacent in  $G$  if they both lie in  $S^{n-1}$  and are antipodal, or if one is a face of the other and  $\sigma \cap \tau$  is  $\text{sgn}(\sigma \cup \tau)$ -labelled. The degree of a completely labelled simplex is the number of completely labelled simplices adjacent to it in  $G$ .

Figure 1 illustrates completely labelled simplices and adjacency for  $n = 2$ .

Proposition.

- (a) The 0-simplex  $\{0\}$  has degree 1;
- (b) Each completely labelled simplex containing a complementary 1-simplex has degree 1;
- (c) Every other completely labelled simplex has degree 2.

Proof. Let  $s$  be the sign vector of the completely labelled simplex  $\sigma$  and suppose  $s$  has  $k$  nonzero components. Then  $\sigma$  lies in the  $k$ -dimensional orthant  $C(s)$ . In addition, the vertices of  $\sigma$  must contain at least  $k$  distinct labels, since  $\sigma$  is  $s$ -labelled. Hence  $\sigma$  is a  $(k-1)$ - or a  $k$ -simplex.

Suppose first that  $\sigma$  is a  $(k-1)$ -simplex. If  $\sigma$  does not lie in  $S^{n-1}$ , it is a face of precisely two  $k$ -simplices in  $C(s)$ , both completely labelled since  $\sigma$  is. If  $\sigma$  lies in  $S^{n-1}$ , it is a face of one completely labelled  $k$ -simplex in  $C(s)$ , and its antipode is completely labelled. In either case,  $\sigma$  is of type (c) and has degree 2.

Suppose now  $\sigma$  is a  $k$ -simplex. It then has  $k+1$  vertices, with one extra label besides the  $k$  it is forced to have by completeness. This other label is either a duplicate of one of the  $k$ , the complement of one of the  $k$ , or  $\pm j$  with  $s_j = 0$ . In the first case,  $\sigma$  has two faces with all  $k$  labels and both are completely labelled;  $\sigma$  is of type (c) and has degree 2. In the second case,  $\sigma$  has just one face with the required  $k$  labels;



$\sigma$  is of type (b) and has degree 1. In the last case, suppose the extra label is  $+j$  ( $-j$ ) and let  $t$  be a sign vector agreeing with  $s$  except that  $t_j = +1$  ( $-1$ ). Then  $\sigma$  is a face of a unique  $(k+1)$ -simplex in  $C(t)$  and this simplex is completely labelled. In addition,  $\sigma$  has one face with the required  $k$  labels; the only exception is when  $\sigma$  is the 0-simplex  $\{0\}$ . Hence  $\sigma$  is either  $\{0\}$  and has degree 1 or is of type (c) with degree 2. The proposition is now proved.

The combinatorial lemma follows directly from the proposition, since every graph has an even number of nodes of odd degree. Indeed, we have a stronger result; there is an odd number of completely labelled simplices containing a complementary 1-simplex. However, since some complementary 1-simplices are contained in no completely labelled simplex and others in several, we can say nothing of the parity of complementary 1-simplices.

More than just a proof, we now have an algorithm: follow a path of adjacent completely labelled simplices from the 0-simplex  $\{0\}$ . By the proposition, the path can terminate only when it encounters a complementary 1-simplex. In Figure 1 the sequence is  $\rho_0, \tau_1, \sigma_2, \dots, \tau_{14}$ . The algorithm can be stated more concisely as follows; this description corresponds to Reiser's algorithm in all respects except the reflection step.

#### Algorithm

Step 0 (Initialization). Set  $s = v = 0 \in R^n$ ,  $\sigma = \{v\}$ . Go to Step 1.

Step 1 (Labelling). Find the label  $\epsilon_i$  of  $v$ ,  $\epsilon = \pm 1$ ,  $1 \leq i \leq n$ . If it is the complement of the label of another vertex of  $\sigma$ , stop. If  $s_i = 0$  go to Step 2, otherwise to Step 3.


Step 2 (Increasing the dimension). Set  $s_i = \varepsilon$ . Let  $\tau$  be the simplex of  $C(s)$  with  $\sigma$  as a face, and  $v$  its new vertex. Set  $\sigma \leftarrow \tau$  and go to Step 1.

Step 3 (Dropping a vertex). Find the vertex  $w \neq v$  of  $\sigma$  with label  $ci$ . Let  $\tau$  be the face of  $\sigma$  opposite  $w$ . If  $\tau$  lies in  $S^{n-1}$ , set  $\sigma \leftarrow -\sigma$ ,  $s \leftarrow -s$ ,  $v \leftarrow -w$  and go to Step 1. If  $\text{sgn}(\tau) = s$ , let  $\rho$  be the simplex in  $C(s)$  with  $\tau$  as a face distinct from  $\sigma$ , and let  $v$  be its new vertex; set  $\sigma \leftarrow \rho$  and go to Step 1. Otherwise go to Step 4.

Step 4 (Decreasing the dimension). Find  $i$  with  $(\text{sgn}(\tau))_i = 0$  and  $s_i \neq 0$ . Set  $\sigma \leftarrow \tau$ ,  $\varepsilon \leftarrow s_i$ ,  $s \leftarrow \text{sgn}(\tau)$ ,  $v \leftarrow w$  and go to Step 3.

In this form the algorithm generates  $\rho_0, \tau_1, \sigma_2, \sigma_4, \sigma_6, \sigma_9, \sigma_{11}, \tau_{12}$  and  $\tau_{14}$  as its successive  $\sigma$ 's in Figure 1.

Note that the only requirement on the labelling is that antipodal vertices in  $S^{n-1}$  have complementary labels--the coordinate structure of  $B^n$  is immaterial. Hence the labels can be permuted so that complementary labels remain complementary and the algorithm run again. This is a possible method for obtaining several complementary 1-simplices--but of course it is not guaranteed to find more than one. In Figure 1, if we interchange labels 1 and -2, -1 and 2 then the completely labelled simplex  $\hat{\sigma}$  containing the complementary 1-simplex  $\hat{\tau}$  is generated.



All simplices marked are completely labelled.  $\sigma$ 's are 2-simplices,  $\tau$ 's 1-simplices and  $\rho$ 's 0-simplices.  $\tau_{14}$  and  $\hat{\tau}$  are complementary. All simplices with consecutive indices are adjacent. Since  $\hat{\tau}$  is not (1,-1)-labelled,  $\hat{\sigma}$  and  $\hat{\tau}$  are not adjacent.

Figure 1

### References

- [1] Barany, I., "Borsuk's Theorem through Complementary Pivoting," manuscript, Budapest, 1977.
- [2] Lefschetz, S., Introduction to Topology, Princeton University Press, Princeton, New Jersey, 1949.
- [3] Meyerson, M., and A. Wright, "A New and Constructive Proof of the Borsuk-Ulam theorem," Proceedings of the A.M.S. 73 (1979), 134-136.
- [4] Reiser, P.M., "A Modified Integer Labeling for Complementarity Algorithms," manuscript, University of Zurich, 1978.
- [5] Todd, M.J., The Computation of Fixed Points and Applications, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [6] Tucker, A.W., "Some topological properties of disk and sphere," Proceedings of the First Canadian Mathematical Congress (Montreal, 1945), 285-309.
- [7] Tucker, A.W., "Antipodal-point Theorems Proved by an Elementary Lemma," Bulletin of the A.M.S. 50 (1944), p. 681.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 SOL-80-12	2. GOVT ACCESSION NO. AD-A090797	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A CONSTRUCTIVE PROOF OF TUCKER'S COMBINATORIAL LEMMA		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) 10 Robert M. Freund & Michael J. Todd		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research - SOL Stanford University Stanford, CA 94305		8. CONTRACT OR GRANT NUMBER(s) DAAG-29-78-G-0026 DOT-DE-1073-76
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NV 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 11 June 1980
		13. NUMBER OF PAGES 8
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  11 75-54.27-10		
18. SUPPLEMENTARY NOTES THE VIEW, OPINIONS, AND/OR FINDINGS CONTAINED IN THIS REPORT ARE THOSE OF THE AUTHOR(S) AND NOT OF THE ARMY. NO OFFICIAL AN OFFICIAL EVALUATION OF THE REPORT HAS BEEN MADE, OR DE- CISION, UNLESS SO DESIGNATED BY OTHER DOCUMENTATION.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  ANTIPODAL POINTS TRIANGULATION COMBINATORICS SIMPLICIAL ALGORITHM		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  Tucker's combinatorial lemma is concerned with certain labellings of the vertices of a triangulation of the n-ball. It can be used as a basis for the proof of antipodal-point theorems in the same way that Sperner's lemma yields Brouwer's theorem. Here we give a constructive proof, which thereby yields algorithms for antipodal-point problems. Our method is based on an algorithm of Reiser.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE  
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

408765